SUCCESSIVE DIFFERENTIATION AND LEIBNITZ'S THEOREM

1.1 Introduction

Successive Differentiation is the process of differentiating a given function successively n times and the results of such differentiation are called successive derivatives. The higher order differential coefficients are of utmost importance in scientific and engineering applications.

Let f(x) be a differentiable function and let its successive derivatives be denoted by $f'(x), f''(x), ..., f^{(n)}(x)$.

Common notations of higher order Derivatives of y = f(x)

1st Derivative:
$$f'(x)$$
 or y' or y_1 or $\frac{dy}{dx}$ or Dy

2nd Derivative:
$$f''(x)$$
 or y'' or y_2 or $\frac{d^2y}{dx^2}$ or D^2y

:

$$n^{th}$$
 Derivative: $f^{(n)}(x)$ or $y^{(n)}$ or y_n or $\frac{d^n y}{dx^n}$ or $D^n y$

1.2 Calculation of nth Derivatives

i. n^{th} Derivative of e^{ax}

Let
$$y = e^{ax}$$

 $y_1 = ae^{ax}$
 $y_2 = a^2e^{ax}$
 \vdots
 $y_n = a^n e^{ax}$

ii. n^{th} Derivative of $(ax + b)^m$, m is a +ve integer greater than n

$$= \frac{m!}{(m-n)!} a^n (ax+b)^{m-n}$$

iii.
$$n^{th}$$
 Derivative of $y = \log(ax + b)$

Let
$$y = \log(ax + b)$$

$$y_1 = \frac{a}{(ax+b)}$$

$$y_2 = \frac{-a^2}{(ax+b)^2}$$

$$y_3 = \frac{2! a^3}{(ax+b)^3}$$

$$\vdots$$

$$y_n = (-1)^{n-1} \frac{(n-1)! a^n}{(ax+b)^n}$$

iv. n^{th} Derivative of $y = \sin(ax + b)$

Let
$$y = \sin(ax + b)$$

$$y_1 = a\cos(ax + b) = a\sin\left(ax + b + \frac{\pi}{2}\right)$$

$$y_2 = a^2\cos\left(ax + b + \frac{\pi}{2}\right) = a^2\sin\left(ax + b + \frac{2\pi}{2}\right)$$

$$\vdots$$

$$y_n = a^n\sin\left(ax + b + \frac{n\pi}{2}\right)$$

Similarly if
$$y = \cos(ax + b)$$

$$y_n = a^n \cos\left(ax + b + \frac{n\pi}{2}\right)$$

v.
$$n^{th}$$
 Derivative of $y = e^{ax} \sin(ax + b)$

Let
$$y = e^{ax}\sin(bx + c)$$

 $y_1 = a e^{ax}\sin(bx + c) + e^{ax}b\cos(bx + c)$
 $= e^{ax} [a\sin(bx + c) + b\cos(bx + c)]$
 $= e^{ax} [r\cos\alpha\sin(bx + c) + r\sin\alpha\cos(bx + c)]$
Putting $a = r\cos\alpha$, $b = r\sin\alpha$

$$= e^{ax} r \sin(bx + c + \alpha)$$
Similarly $y_2 = e^{ax} r^2 \sin(bx + c + 2\alpha)$
:

$$y_n = e^{ax} r^n \sin(bx + c + n\alpha)$$
where $r^2 = a^2 + b^2$ and $\tan \alpha = \frac{b}{a}$

$$\therefore y_n = e^{ax} (a^2 + b^2)^{\frac{n}{2}} \sin(bx + c + n \tan^{-1} \frac{b}{a})$$

Similarly if
$$y = e^{ax}\cos(ax + b)$$

$$y_n = e^{ax} r^n \cos(bx + c + n\alpha)$$

= $e^{ax} (a^2 + b^2)^{\frac{n}{2}} \cos(bx + c + n \tan^{-1} \frac{b}{a})$

Summary of Results

Function	n^{th} Derivative
$y = e^{ax}$	$y_n = a^n e^{ax}$
$y = (ax + b)^m$	$y_n = \begin{cases} \frac{m!}{(m-n)!} a^n (ax+b)^{m-n}, m > 0, m > n \\ 0, & m > 0, & m < n, \\ n! \ a^n, & m = n \\ \frac{(-1)^n n! \ a^n}{(ax+b)^{n+1}}, & m = -1 \end{cases}$
$y = \log(ax + b)$	$y_n = (-1)^{n-1} \frac{(n-1)! \ a^n}{(ax+b)^n}$
	$y_n = a^n \sin\left(ax + b + \frac{n\pi}{2}\right)$
y = cos(ax + b)	$y_n = a^n \cos\left(ax + b + \frac{n\pi}{2}\right)$
$y = e^{ax} \sin(bx + c)$	$y_n = e^{ax} (a^2 + b^2)^{\frac{n}{2}} \sin(bx + c + n \tan^{-1} \frac{b}{a})$
$y = e^{ax} \cos(bx + c)$	$y_n = e^{ax} (a^2 + b^2)^{\frac{n}{2}} \cos(bx + c + n \tan^{-1} \frac{b}{a})$

Example 1 Find the n^{th} derivative of $\frac{1}{1-5x+6x^2}$

Solution: Let
$$y = \frac{1}{1 - 5x + 6x^2}$$

Resolving into partial fractions

$$y = \frac{1}{1 - 5x + 6x^2} = \frac{1}{(1 - 3x)(1 - 2x)} = \frac{3}{1 - 3x} - \frac{2}{1 - 2x}$$

$$\therefore y_n = \frac{3(-3)^n (-1)^n n!}{(1 - 3x)^{n+1}} - \frac{2(-2)^n (-1)^n n!}{(1 - 2x)^{n+1}}$$

$$\Rightarrow y_n = (-1)^{n+1} n! \left[\left(\frac{3}{1 - 3x} \right)^{n+1} - \left(\frac{2}{1 - 2x} \right)^{n+1} \right]$$

Example 2 Find the n^{th} derivative of $\sin 6x \cos 4x$

Solution: Let
$$y = \sin 6x \cos 4x$$

$$= \frac{1}{2} (\sin 10 x + \cos 2 x)$$

$$\therefore y_n = \frac{1}{2} \left[10^n \sin \left(10x + \frac{n\pi}{2} \right) + 2^n \cos \left(2x + \frac{n\pi}{2} \right) \right]$$

Example 3 Find n^{th} derivative of sin^2xcos^3x

Solution: Let
$$y = sin^2 x cos^3 x$$

$$= \sin^{2}x\cos^{2}x \cos x$$

$$= \frac{1}{4}\sin^{2}2x \cos x = \frac{1}{8}(1 - \cos 4x)\cos x$$

$$= \frac{1}{8}\cos x - \frac{1}{8}\cos 4x \cos x$$

$$= \frac{1}{8}\cos x - \frac{1}{16}(\cos 3x + \cos 5x)$$

$$= \frac{1}{16}(2\cos x - \cos 3x - \cos 5x)$$

$$\therefore y_{n} = \frac{1}{16}\left[2\cos\left(x + \frac{n\pi}{2}\right) - 3^{n}\cos\left(3x + \frac{n\pi}{2}\right) - 5^{n}\cos\left(5x + \frac{n\pi}{2}\right)\right]$$

Example 4 Find the n^{th} derivative of sin^4x

Solution: Let
$$y = \sin^4 x = (\sin^2 x)^2$$

$$= \left(\frac{1}{2} 2 \sin^2 x\right)^2$$

$$= \frac{1}{4} \left((1 - \cos 2x)^2 \right)$$

$$= \frac{1}{4} \left[1 - 2\cos 2x + \frac{1}{2} (2\cos^2 2x) \right]$$

$$= \frac{1}{4} \left[1 - 2\cos 2x + \frac{1}{2} (1 + \cos 4x) \right]$$

$$= \frac{3}{8} - \frac{1}{2}\cos 2x + \frac{1}{8}\cos 4x$$

$$\therefore y_n = -\frac{1}{2} 2^n \cos \left(2x + \frac{n\pi}{2} \right) + \frac{1}{8} 4^n \cos \left(4x + \frac{n\pi}{2} \right)$$

Example 5 Find the n^{th} derivative of $e^{3x}\cos x \sin^2 2x$

Solution: Let $y = e^{3x} \cos x \sin^2 2x$

Example 6 If $y = \sin ax + \cos ax$, prove that $y_n = a^n [1 + (-1)^n \sin 2ax]^{\frac{1}{2}}$ **Solution:** $y = \sin ax + \cos ax$

$$\therefore y_n = a^n \left[\sin \left(ax + \frac{n\pi}{2} \right) + \cos \left(ax + \frac{n\pi}{2} \right) \right]$$

$$= a^{n} \left[\left\{ \sin \left(ax + \frac{n\pi}{2} \right) + \cos \left(ax + \frac{n\pi}{2} \right) \right\}^{2} \right]^{\frac{1}{2}}$$

$$= a^{n} \left[\sin^{2} \left(ax + \frac{n\pi}{2} \right) + \cos^{2} \left(ax + \frac{n\pi}{2} \right) + 2 \sin \left(ax + \frac{n\pi}{2} \right) \cdot \cos \left(ax + \frac{n\pi}{2} \right) \right]^{\frac{1}{2}}$$

$$= a^{n} \left[1 + \sin (2ax + n\pi) \right]^{\frac{1}{2}}$$

$$= a^{n} \left[1 + \sin 2ax \cos n\pi + \cos 2ax \sin n\pi \right]^{\frac{1}{2}}$$

$$= a^{n} \left[1 + (-1)^{n} \sin 2ax \right]^{\frac{1}{2}} \quad \because \cos n\pi = (-1)^{n} \text{ and } \sin n\pi = 0$$

Example 7 Find the n^{th} derivative of $\tan^{-1} \frac{x}{a}$

Solution: Let
$$y = \tan^{-1} \frac{x}{a}$$

$$\Rightarrow y_1 = \frac{dy}{dx} = \frac{1}{a\left(1 + \frac{x^2}{a^2}\right)} = \frac{a}{x^2 + a^2} = \frac{a}{x^2 - (ai)^2}$$

$$= \frac{a}{(x + ai)(x - ai)} = \frac{a}{2ai} \left(\frac{1}{x - ai} - \frac{1}{x + ai}\right)$$

$$= \frac{1}{2i} \left(\frac{1}{x - ai} - \frac{1}{x + ai}\right)$$

Differentiating above (n-1) times w.r.t. x, we get

$$y_n = \frac{1}{2i} \left[\frac{(-1)^{n-1}(n-1)!}{(x-ai)^n} - \frac{(-1)^{n-1}(n-1)!}{(x+ai)^n} \right]$$

Substituting $x = r \cos\theta$, $a = r \sin\theta$ such that $\theta = \tan^{-1} \frac{x}{a}$

$$\Rightarrow y_n = \frac{(-1)^{n-1}(n-1)!}{2i} \left[\frac{1}{r^n(\cos\theta - i\sin\theta)^n} - \frac{1}{r^n(\cos\theta + i\sin\theta)^n} \right]$$
$$= \frac{(-1)^{n-1}(n-1)!}{2ir^n} \left[(\cos\theta - i\sin\theta)^{-n} - (\cos\theta + i\sin\theta)^{-n} \right]$$

Using De Moivre's theorem, we get

$$y_n = \frac{(-1)^{n-1}(n-1)!}{2ir^n} \left[\cos n\theta + i \sin n\theta - \cos n\theta + i \sin n\theta \right]$$

$$= \frac{(-1)^{n-1}(n-1)!}{r^n} \sin n\theta$$

$$= \frac{(-1)^{n-1}(n-1)!}{\left(\frac{a}{\sin \theta}\right)^n} \sin n\theta \qquad \because a = r \sin \theta$$

$$= \frac{(-1)^{n-1}(n-1)!}{a^n} \sin n\theta \sin n\theta \qquad \text{where } \theta = \tan^{-1}\frac{a}{x}$$

Example 8 Find the n^{th} derivative of $\frac{1}{1+x+x^2}$

Solution: Let
$$y = \frac{1}{1+x+x^2}$$

$$= \frac{1}{(x-w)(x-w^2)} \text{ where } w = \frac{-1+i\sqrt{3}}{2} \text{ and } w^2 = \frac{-1-i\sqrt{3}}{2}$$
Resolving into partial fractions
$$y = \frac{1}{w-w^2} \left(\frac{1}{x-w} - \frac{1}{x-w^2} \right)$$

$$= \frac{1}{i\sqrt{3}} \left(\frac{1}{x-w} - \frac{1}{x-w^2} \right) = \frac{-i}{\sqrt{3}} \left(\frac{1}{x-w} - \frac{1}{x-w^2} \right)$$

Differentiating n times w.r.t., we get

$$\begin{split} y_n &= \frac{-i}{\sqrt{3}} \left[\frac{(-1)^n n!}{(x-w)^{n+1}} - \frac{(-1)^n n!}{(x-w^2)^{n+1}} \right] \\ &= \frac{-i \, (-1)^n n!}{\sqrt{3}} \left[\frac{1}{(x-w)^{n+1}} - \frac{1}{(x-w^2)^{n+1}} \right] \\ &= \frac{i \, (-1)^{n+1} n!}{\sqrt{3}} \left[\frac{1}{\left(x + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^{n+1}} - \frac{1}{\left(x + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^{n+1}} \right] \\ &= \frac{i \, 2^{n+1} \, (-1)^{n+1} n!}{\sqrt{3}} \left[\frac{1}{(2x+1-i\sqrt{3})^{n+1}} - \frac{1}{(2x+1+i\sqrt{3})^{n+1}} \right] \end{split}$$

Substituting $2x + 1 = r \cos\theta$, $\sqrt{3} = r \sin\theta$ such that $\theta = \tan^{-1} \frac{\sqrt{3}}{2x+1}$ $y_n = \frac{i \, 2^{n+1} \, (-1)^{n+1} n!}{\sqrt{3} \, r^{n+1}} \left[(\cos\theta - i \sin\theta)^{-(n+1)} - (\cos\theta + i \sin\theta)^{-(n+1)} \right]$

Using De Moivre's theorem, we get

$$y_n = \frac{i \, 2^{n+1} \, (-1)^{n+1} n!}{\sqrt{3} \, \left(\frac{\sqrt{3}}{\sin \theta}\right)^{n+1}} \left[\cos(n+1)\theta + i \sin(n+1)\theta - \cos(n+1)\theta + i \sin(n+1)\theta \right]$$

$$\because \sqrt{3} = r \sin\theta$$

$$= \frac{i \, 2^{n+1} \, (-1)^{n+1} n!}{\left(\sqrt{3}\right)^{n+2}} \, 2i \, \sin(n+1)\theta \, \sin^{n+1}\theta$$

$$= \frac{(-2)^{n+2} n!}{\left(\sqrt{3}\right)^{n+2}} \, \sin(n+1)\theta \, \sin^{n+1}\theta \, \cos^{n+1}\theta$$

$$= \frac{(-2)^{n+2} n!}{\sqrt{3}^{n+2}} \sin(n+1)\theta \sin^{n+1}\theta \quad \text{where } \theta = \tan^{-1} \frac{\sqrt{3}}{2x+1}$$

Example 9 If $y = x + \tan x$, show that $\cos^2 x \frac{d^2 y}{dx^2} - 2y + 2x = 0$

Solution: $y = x + \tan x$

$$\Rightarrow \frac{\mathrm{d}y}{\mathrm{d}x} = 1 + \sec^2 x$$

$$\frac{d^2y}{dx^2} = 2 \sec x (\sec x \tan x) = 2 \sec^2 x \tan x$$

Example 10 If $y = \log(x + \sqrt{x^2 + 1})$, show that $(1 + x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 0$

Solution: $y = \log(x + \sqrt{x^2 + 1})$

$$\Rightarrow \frac{\mathrm{dy}}{\mathrm{dx}} = \frac{1 + \frac{x}{\sqrt{1 + x^2}}}{x + \sqrt{1 + x^2}} = \frac{1}{\sqrt{1 + x^2}}$$

$$\Rightarrow (\sqrt{1+x^2}) \frac{\mathrm{d}y}{\mathrm{d}x} = 1$$

Differentiating both sides w.r.t. x, we get

$$(\sqrt{1+x^2}) \frac{d^2 y}{dx^2} + \frac{x}{\sqrt{1+x^2}} \frac{dy}{dx} = 0$$

$$\Rightarrow (1+x^2) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = 0$$

Exercise 1 A

1. Find the n^{th} derivative of $\frac{x^4}{(x-1)(x-2)}$

Ans.
$$(-1)^n n! \left[\frac{16}{(x-2)^{n+1}} - \frac{1}{(x-1)^{n+1}} \right]$$

2. Find the n^{th} derivative of $\cos x \cos 2x \cos 3x$

$$\operatorname{Ans.} \frac{1}{4} \left[2^n \cos \left(2x + \frac{n\pi}{2} \right) + 4^n \cos \left(4x + \frac{n\pi}{2} \right) + 6^n \cos \left(6x + \frac{n\pi}{2} \right) \right]$$

- 3. If $x = \sin t$, $y = \sin at$, show that $(1 x^2) \frac{d^2y}{dx^2} x \frac{dy}{dx} + a^2y = 0$
- 4. If $p^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta$, show that $p + \frac{d^2 p}{d\theta^2} = \frac{a^2 b^2}{p^3}$
- 5. If $y = \frac{x}{x^2 + a^2}$, find y_n i.e. the n^{th} derivative of y

Ans.
$$\frac{(-1)^n n!}{a^{n+1}} \cos(n+1)\theta \sin^{n+1}\theta$$
 where $\theta = \tan^{-1} \frac{x}{a}$

6. If $y = e^x \sin^2 x$, find y_n i.e. the n^{th} derivative of y

Ans.
$$\frac{1}{2} e^x \left[1 - 16 \left(2x + \frac{n\pi}{2} \right) \right]$$

7. Find n^{th} differential coefficient of $y = \log[(ax + b)(cx + d)]$

Ans.
$$y_n = (-1)^{n-1}(n-1)! \left[\frac{a^n}{(ax+b)^n} + \frac{c^n}{(cx+d)^n} \right]$$

- 8. If $y = x \log \frac{x-1}{x+1}$, show that $y_n = (-1)^{n-1}(n-2)! \left[\frac{x-n}{(x-1)^n} + \frac{x+n}{(x+1)^n} \right]$
- 9. If $y = \tan^{-1} \frac{\sqrt{1+x^2}-1}{x}$, show that $y_n = \frac{1}{2}(-1)^{n-1}(n-1)! \sin n\theta \sin^n\theta$

1.2 LEIBNITZ'S THEOREM

If u and v are functions of x such that their n^{th} derivatives exist, then the n^{th} derivative of their product is given by

 $(u\ v)_n = u_n v + n_{C_1} u_{n-1} v_1 + n_{C_2} u_{n-2} v_2 + \dots + n_{C_r} u_{n-r} v_r + \dots + uv_n$ where u_r and v_r represent r^{th} derivatives of u and v respectively.

Example11 Find the n^{th} derivative of $x \log x$

Solution: Let $u = \log x$ and v = x

Then
$$u_n = (-1)^{n-1} \frac{(n-1)!}{x^n}$$
 and $u_{n-1} = (-1)^{n-2} \frac{(n-2)!}{x^{n-1}}$

By Leibnitz's theorem, we have

By Lefthitz's theorem, we have
$$(u v)_n = u_n v + n_{C_1} u_{n-1} v_1 + n_{C_2} u_{n-2} v_2 + \dots + n_{C_r} u_{n-r} v_r + \dots + u v_n$$

$$\Rightarrow (x \log x)_n = (-1)^{n-1} \frac{(n-1)!}{x^n} x + n(-1)^{n-2} \frac{(n-2)!}{x^{n-1}} + 0$$

$$= (-1)^{n-1} \frac{(n-1)!}{x^{n-1}} + n(-1)^{n-2} \frac{(n-2)!}{x^{n-1}}$$

$$= (-1)^{n-2} \frac{(n-2)!}{x^{n-1}} [-(n-1) + n]$$

$$= (-1)^{n-2} \frac{(n-2)!}{x^{n-1}}$$

Example 12 Find the n^{th} derivative of $x^2e^{3x} \sin 4x$

Solution: Let $u = e^{3x} \sin 4x$ and $v = x^2$

Then
$$u_n = e^{3x} 25^{\frac{n}{2}} \sin\left(4x + n \tan^{-1} \frac{4}{3}\right)$$

= $e^{3x} 5^n \sin\left(4x + n \tan^{-1} \frac{4}{3}\right)$

By Leibnitz's theorem, we have

$$(u v)_n = u_n v + n_{C_1} u_{n-1} v_1 + n_{C_2} u_{n-2} v_2 + \dots + n_{C_r} u_{n-r} v_r + \dots + u v_n$$

$$\Rightarrow \left(x^2 e^{3x} \sin 4x \right)_n = x^2 e^{3x} 5^n \sin \left(4x + n \tan^{-1} \frac{4}{3} \right) +$$

$$2nx e^{3x} 5^{n-1} \sin \left(4x + (n-1) \tan^{-1} \frac{4}{3} \right) +$$

$$n(n-1) e^{3x} 5^{n-2} \sin \left(4x + (n-2) \tan^{-1} \frac{4}{3} \right) + 0$$

$$= e^{3x} 5^n \left[x^2 \sin\left(4x + n \tan^{-1}\frac{4}{3}\right) + \frac{2nx}{5} \sin\left(4x + (n-1)\tan^{-1}\frac{4}{3}\right) + \frac{n(n-1)}{25} \sin\left(4x + (n-2)\tan^{-1}\frac{4}{3}\right) \right]$$

Example 13 If $y = a \cos(\log x) + b \sin(\log x)$, show that

$$x^{2}y_{n+2} + (2n+1)xy_{n+1} + n(n+1)y_{n} = 0$$

Solution: Here
$$y = a \cos(\log x) + b \sin(\log x)$$

$$\Rightarrow y_1 = \frac{-a}{x}\sin(\log x) + \frac{b}{x}\cos(\log x)$$

$$\Rightarrow xy_1 = -a\sin(\log x) + b\cos(\log x)$$

Differentiating both sides w.r.t. x, we get

$$xy_2 + y_1 = -\frac{a}{x}\cos(\log x) + \frac{-b}{x}\sin(\log x)$$

$$\Rightarrow x^2y_2 + xy_1 = -\{a\cos(\log x) + b\sin(\log x)\}$$

$$= -y$$

$$\Rightarrow x^2y_2 + xy_1 + y = 0$$

Using Leibnitz's theorem, we get

$$(y_{n+2}x^2 + n_{C_1}y_{n+1}2x + n_{C_2}y_n.2) + (y_{n+1}x + n_{C_1}y_n.1) + y_n = 0$$

$$\Rightarrow y_{n+2}x^2 + y_{n+1}2nx + n(n-1)y_n + y_{n+1}x + ny_n + y_n = 0$$

$$\Rightarrow x^2y_{n+2} + (2n+1)xy_{n+1} + (n^2+1)y_n = 0$$

Example 14 If $y = \log (x + \sqrt{1 + x^2})$

Prove that
$$(1 + x^2)y_{n+2} + (2n + 1)xy_{n+1} + n^2y_n = 0$$

Solution:
$$y = \log (x + \sqrt{1 + x^2})$$

$$\Rightarrow y_1 = \frac{1}{x + \sqrt{1 + x^2}} \left(1 + \frac{1}{2\sqrt{1 + x^2}} 2x \right) = \frac{1}{\sqrt{1 + x^2}}$$

$$\Rightarrow (1 + x^2) y_1^2 = 1$$

Differentiating both sides w.r.t. x, we get

$$(1 + x2)2y1y2 + 2xy12 = 0$$

$$\Rightarrow (1 + x2)y2 + xy1 = 0$$

$$[y_{n+2}(1+x^2) + n_{C_1}y_{n+1}2x + n_{C_2}y_n.2] + (y_{n+1}x + n_{C_1}y_n.1) = 0$$

$$\Rightarrow y_{n+2}(1+x^2) + y_{n+1}2nx + n(n-1)y_n + y_{n+1}x + ny_n = 0$$

$$\Rightarrow (1+x^2)y_{n+2} + (2n+1)xy_{n+1} + n^2y_n = 0$$

Example 15 If $y = \sin(m \sin^{-1} x)$, show that

$$(1-x^2)y_{n+2} = (2n+1)xy_{n+1} + (n^2-m^2)y_n$$
. Also find $y_n(0)$

Solution: Here $y = \sin(m \sin^{-1} x)$ ①

$$\Rightarrow y_1 = \frac{m}{\sqrt{1-x^2}} \cos(m \sin^{-1} x) \qquad \dots 2$$

$$\Rightarrow (1 - x^2)y_1^2 = m^2\cos^2(m \sin^{-1}x)$$

$$\Rightarrow (1 - x^2)y_1^2 = m^2[1 - \sin^2(m \sin^{-1}x)]$$

$$\Rightarrow (1 - x^2)y_1^2 = m^2(1 - y^2).....$$

$$\Rightarrow$$
 $(1-x^2)y_1^2 + m^2y^2 = m^2$

Differentiating w.r.t. x, we get

$$(1 - x^2)2y_1y_2 + y_1^2(-2x) + m^22yy_1 = 0$$

$$\Rightarrow (1 - x^2)y_2 - xy_1 + m^2y = 0$$

Using Leibnitz's theorem, we get

$$[y_{n+2}(1-x^2) + n_{c_1}y_{n+1}(-2x) + n_{c_2}y_n(-2)] - (y_{n+1}x + n_{c_1}y_n1) + m^2y_n = 0$$

$$\Rightarrow y_{n+2}(1-x^2) - y_{n+1}2nx - n(n-1)y_n - (y_{n+1}x + ny_n) + m^2y_n = 0$$

$$\Rightarrow (1 - x^2)y_{n+2} = (2n + 1)xy_{n+1} + (n^2 - m^2)y_n \dots 4$$

Putting x = 0 in ①, ②and ③

$$y(0) = 0, y_1(0) = m \text{ and } y_2(0) = 0$$

Putting x = 0 in 4

$$y_{n+2}(0) = (n^2 - m^2)y_n(0)$$

Putting $n = 1,2,3 \dots$ in the above equation, we get

$$y_3(0) = (1^2 - m^2)y_1(0)$$

$$=(1^2-m^2)m$$
 $y_1(0)=m$

$$y_4(0) = (2^2 - m^2)y_2(0)$$

$$= 0 y_2(0) = 0$$

$$y_5(0) = (3^2 - m^2)y_3(0)$$

$$= m(1^2 - m^2)(3^2 - m^2)$$
.

:

$$\Rightarrow y_n(0) = \begin{cases} 0, & \text{if n is even} \\ m(1^2 - m^2)(3^2 - m^2) \dots [(n-2)^2) - m^2 \end{bmatrix}, \text{if n is odd} \end{cases}$$

Example 16 If
$$y = e^{msin^{-1}x}$$
, show that $(1-x^2)y_{n+2} - (2n+1)xy_{n+1} - (n^2+m^2)y_n = 0$. Also find $y_n(0)$.

Solution: Here
$$y = e^{msin^{-1}x}$$
 ...①

$$\Rightarrow y_1 = \frac{m}{\sqrt{1-x^2}} e^{m \sin^{-1} x}$$
$$= \frac{my}{\sqrt{1-x^2}} \dots (2)$$

$$\Rightarrow (1 - x^2)y_1^2 = m^2y^2$$

Differentiating above equation w.r.t. x, we get

$$(1 - x^2)2y_1y_2 + y_1^2(-2x) = m^2 2yy_1$$

$$\Rightarrow (1 - x^2)y_2 - xy_1 - m^2y = 0 \qquad \dots \dots 3$$

Differentiating above equation n times w.r.t. x using Leibnitz's theorem, we get

$$[y_{n+2}(1-x^2) + n_{C_1}y_{n+1}(-2x) + n_{C_2}y_n(-2)] - (y_{n+1}x + n_{C_1}y_n1) - m^2y_n = 0$$

$$\Rightarrow y_{n+2}(1-x^2) - y_{n+1}2nx - n(n-1)y_n - (y_{n+1}x + ny_n) - m^2y_n = 0$$

$$\Rightarrow (1-x^2)y_{n+2} - (2n+1)xy_{n+1} - (n^2+m^2)y_n = 0.....4$$

To find $y_n(0)$: Putting x = 0 in ①, ② and ③

$$y(0) = 1$$
, $y_1(0) = m$ and $y_2(0) = m^2$

Also putting x = 0 in , we get

$$y_{n+2}(0) = (n^2 + m^2)y_n(0)$$

Putting n = 1,2,3 ... in the above equation, we get

$$y_{3}(0) = (1^{2} + m^{2})y_{1}(0)$$

$$= (1^{2} + m^{2})m \qquad \because y_{1}(0) = m$$

$$y_{4}(0) = (2^{2} + m^{2})y_{2}(0)$$

$$= m^{2}(2^{2} + m^{2}) \qquad \because y_{2}(0) = m^{2}$$

$$y_{5}(0) = (3^{2} + m^{2})y_{3}(0)$$

$$= m(1^{2} + m^{2})(3^{2} + m^{2})$$

$$\vdots$$

$$\Rightarrow y_n(0) = \begin{cases} m^2(2^2 + m^2) \dots [(n-2)^2 + m^2], & \text{if } n \text{ is even} \\ m(1^2 + m^2)(3^2 + m^2) \dots [(n-2)^2 + m^2], & \text{if } n \text{ is odd} \end{cases}$$

Example 17 If $y = tan^{-1}x$, show that

$$(1-x^2)y_{n+2} + 2(n+1)xy_{n+1} + n(n+1)y_n = 0$$
. Also find $y_n(0)$

Solution: Here $y = tan^{-1}x...$

$$\Rightarrow y_1 = \frac{1}{1+x^2} \dots 2$$

$$y_2 = \frac{-2x}{1+x^2}$$

$$\Rightarrow (1+x^2)y_2 + 2xy_1 = 0 \dots 3$$

$$[y_{n+2}(1+x^2) + n_{C_1}y_{n+1}(2x) + n_{C_2}y_n(2)] + 2(y_{n+1}x + n_{C_1}y_n1) = 0$$

$$\Rightarrow y_{n+2}(1+x^2) + y_{n+1}2nx + n(n-1)y_n + 2(y_{n+1}x + ny_n) = 0$$

$$\Rightarrow (1+x^2)y_{n+2} + 2(n+1)xy_{n+1} + n(n+1)y_n = 0......4$$

To find $y_n(0)$: Putting x = 0 in ①, ②and ③, we get

$$y(0) = 0$$
, $y_1(0) = 1$ and $y_2(0) = 0$

Also putting x = 0 in 4, we get

$$y_{n+2}(0) = -n(n+1)y_n(0)$$

Putting n = 1,2,3 ... in the above equation, we get

$$y_{3}(0) = -1(2)y_{1}(0)$$

$$= -2 \qquad \because y_{1}(0) = 1$$

$$y_{4}(0) = -2(3)y_{2}(0)$$

$$= 0 \qquad \because y_{2}(0) = 0$$

$$y_{5}(0) = -3(4)y_{3}(0)$$

$$= -3(4)(-2) = 4!$$

$$y_{6}(0) = -4(5)y_{4}(0) = 0$$

$$y_{7}(0) = -5(6)y_{5}(0) = -5(6)4! = -(6!)$$

$$\vdots$$

$$\Rightarrow y_{2n+1}(0) = (-1)^{n}(2n)! \text{ and } y_{2n}(0) = 0$$
Example 18 If $y = (sin^{-1}x)^{2}$, show that $(1 - x^{2})y_{n+2} - (2n + 1)xy_{n+1} - n^{2}y_{n} = 0$. Also find $y_{n}(0)$
Solution: Here $y = (sin^{-1}x)^{2}$①
$$\Rightarrow y_{1} = 2sin^{-1}x \cdot \frac{1}{\sqrt{1-x^{2}}} \qquad \dots \dots \text{②}$$
Squaring both the sides, we get
$$(1 - x^{2})y_{1}^{2} = 4(sin^{-1}x)^{2}$$

$$\Rightarrow (1 - x^{2})y_{1}^{2} = 4(y)^{2}$$

Differentiating the above equation w.r.t. *x*, we get

$$(1 - x^2)2y_1y_2 + y_1^2(-2x) - 4y_1 = 0$$

$$\Rightarrow (1 - x^2)y_2 + y_1(-x) - 2 = 0 \qquad \dots \dots 3$$

Differentiating the above equation n times w.r.t. x using Leibnitz's theorem, we get

$$[y_{n+2}(1-x^2) + n_{C_1}y_{n+1}(-2x) + n_{C_2}y_n(-2)] - (y_{n+1}x + n_{C_1}y_n1) = 0$$

$$\Rightarrow y_{n+2}(1-x^2) - y_{n+1}2nx - n(n-1)y_n - (y_{n+1}x + ny_n) = 0$$

$$\Rightarrow (1-x^2)y_{n+2} - (2n+1)xy_{n+1} - y_nn^2 = 0......4$$
To find $y_n(0)$: Putting $x = 0$ in ①, ②and ③, we get
$$y(0) = 0, y_1(0) = 0 \text{ and } y_2(0) = 2$$

Also putting x = 0 in 4, we get

$$y_{n+2}(0) = n^2 y_n(0)$$

Putting n = 1,2,3... in the above equation, we get

$$\Rightarrow y_n(0) = \begin{cases} 0, & \text{if } n \text{ is odd} \\ 2 \cdot 2^2 \cdot 4^2 \dots \dots (n-2)^2, & \text{if } n \text{ is even} \end{cases}$$

Exercise 1 B

1 .Find y_n , if $y = x^3 \cos x$

Ans.
$$x^3 cos\left(x + \frac{n\pi}{2}\right) + 3nx^2 cos\left[x + \frac{1}{2}(n-1)\pi\right] + 3n(n-1)xcos\left[x + \frac{1}{2}(n-2)\pi\right] + n(n-1)(n-2)cos\left[x + \frac{1}{2}(n-3)\pi\right]$$

2. Find y_n , if $y = x^2 e^x \cos x$

Ans.
$$2^{\frac{n}{2}}e^x \cos\left(x + \frac{n\pi}{4}\right)$$

3. If
$$y^{\frac{1}{m}} + y^{\frac{-1}{m}} = 2x$$
, prove that $(x^2 - 1)y_{n+2} + (2n+1)xy_{n+1} + (n^2 - m^2)y_n = 0$

4. If
$$y\sqrt{1+x^2} = \log(x+\sqrt{1+x^2})$$
, prove that
$$(1+x^2)y_{n+2} + (2n+3)xy_{n+1} + (n+1)^2y_n = 0$$

5. If
$$y = [x + \sqrt{1 + x^2}]^m$$
, prove that $(x^2 + 1)y_{n+2} + (2n + 1)xy_{n+1} + (n^2 - m^2)y_n = 0$

6 If
$$y = (sinh^{-1}x)^2$$
, show that
$$(1+x^2)y_{n+2} + (2n+1)xy_{n+1} + n^2y_n = 0$$
. Also find $y_n(0)$.

Ans.
$$y_{2n+1}(0) = 0$$
 and $y_{2n}(0) = (-1)^{n-1}2 \cdot 2^2 \cdot 4^2 \cdot \dots \cdot (2n-2)^2$

7. If $y = \cos(m \sin^{-1} x)$, show that $(1 - x^2)y_{n+2} = (2n + 1)xy_{n+1} + (n^2 - m^2)y_n$. Also find $y_n(0)$.

8. If $f(x) = \tan x$, prove that $f^n(0) - n_{c_2} f^{n-2}(0) + n_{c_4} f^{n-4}(0) - \dots = \sin \frac{n\pi}{2}$